

# Dosing for Infants and Children

From Your Healthcare Professional

Concentrated  
**Motrin**  
Infants' Drops

Children's  
**Motrin**

**DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**

If possible, use weight to dose; otherwise use age.

mL = milliliter  
tsp = teaspoon



## Infants' MOTRIN® Concentrated Drops

Active Ingredient:  
Ibuprofen 50 mg (NSAID)<sup>†</sup>  
(in each 1.25 mL)

<sup>†</sup>Nonsteroidal anti-inflammatory drug  
Use only as directed.

Available in:  
 Dye-Free Berry



## Children's MOTRIN® Oral Suspension

Active Ingredient:  
Ibuprofen 100 mg (NSAID)<sup>†</sup>  
(in each 5 mL or 1 tsp)

<sup>†</sup>Nonsteroidal anti-inflammatory drug  
Use only as directed.

Available in:  
 Original Berry  
 Dye-Free Berry

WEIGHT	AGE	Infants' MOTRIN® Concentrated Drops	Children's MOTRIN® Oral Suspension
6-11 lbs	0-5 mos	<b>Do not use</b>	—
12-17 lbs	6-11 mos	<b>1.25 mL</b> 	—
18-23 lbs	12-23 mos	<b>1.875 mL</b> 	—
24-35 lbs	2-3 yrs	—	<b>5 mL</b> (1 tsp)
36-47 lbs	4-5 yrs	—	<b>7.5 mL</b> (1½ tsp)
48-59 lbs	6-8 yrs	—	<b>10 mL</b> (2 tsp)
60-71 lbs	9-10 yrs	—	<b>12.5 mL</b> (2½ tsp)
72-95 lbs	11 yrs	—	<b>15 mL</b> (3 tsp)

## IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: \_\_\_\_\_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products.
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours.
- Do NOT administer longer than 10 days, unless directed by a doctor.
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID<sup>†</sup> (prescription or non-prescription).  
— MOTRIN® contains ibuprofen
- Use only the dosing device that comes with a specific product.

<sup>†</sup>Nonsteroidal anti-inflammatory drug